

AD-A189 178

A RELATIONSHIP BETWEEN OCEAN CIRCULATION AND VOLUME
REGENERATION IN THE (U) NAVAL UNDERWATER SYSTEMS
CENTER NEW LONDON CT NEW LONDON LAB

1/1

UNCLASSIFIED


D G BROWNING ET AL. 26 JUN 86 NUSC-ID-7681 F/G 8/3

ML

END

DATE


8



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963-A

NUSC Technical Document 7681
26 June 1986

AD-A189 178 COPY



A Relationship Between Ocean Circulation and Volume Reverberation in the Subarctic Northeast Pacific Ocean (Gulf of Alaska)

**A Paper Presented at the
111th Meeting of the Acoustical Society of America,
Cleveland, Ohio, 12-16 May 1986**

**David G. Browning
Surface Ship Sonar Department**

**R. G. Turner
J. W. Power
Defence Research Establishment Pacific**

**DTIC
ELECTE
OCT 07 1987
S
D**



**Naval Underwater Systems Center
Newport, Rhode Island / New London, Connecticut**


Approved for public release; distribution unlimited.

37 10 1 332

PREFACE

This research was conducted under NUSC Project No. A60080; Subproject No. RS-11-1220; Principal Investigator, Dr. Azizul H. Quazi (Code 3315); Program Manager, Ronald Tompkins (Code 33A); Sponsor Theo Koj, Office of Naval Technology.

REVIEWED AND APPROVED: 26 June 1986



LARRY FREEMAN
HEAD, SURFACE SHIP SONAR DEPARTMENT

The author of this document is located at the New London Laboratory,
Naval Underwater Systems Center, New London, Connecticut 06320.

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NUSC TD 7681			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Naval Underwater Systems Center		6b. OFFICE SYMBOL (If applicable) 3331	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code). New London Laboratory New London, CT 06320			7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Technology		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
8c. ADDRESS (City, State, and ZIP Code) Washington, DC 20375			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO. A60080	TASK NO.
			WORK UNIT ACCESSION NO.		
11. TITLE (Include Security Classification) A RELATIONSHIP BETWEEN OCEAN CIRCULATION AND VOLUME REVERBERATION IN THE SUBARCTIC NORTHEAST PACIFIC OCEAN (GULF OF ALASKA)					
12. PERSONAL AUTHOR(S) David G. Browning, R. G. Turner, and J. W. Power					
13a. TYPE OF REPORT		13b. TIME COVERED FROM TO		14. DATE OF REPORT (Year, Month, Day) 86-06-26	
				15. PAGE COUNT 14	
16. SUPPLEMENTARY NOTATION A Paper Presented at the 111th Meeting of the Acoustical Society of America, Cleveland, Ohio, 12-16 May 1986					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Ocean Circulation		
			Subarctic Pacific Ocean. <i>✓</i>		
			Volume Reverberation <i>acoustic</i>		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Earlier investigations have shown a significant change in integrated scattering and spectral characteristics when transmitting into the subarctic (above 40 north latitude) northeast Pacific Ocean. An analysis of an extensive series of volume reverberation measurements obtained by Turner indicates a strong influence of the counterclockwise circulation around the Alaskan Gyre on the distribution of scattering strengths. At higher frequencies (5-20 kHz) the greater scattering strengths are found in the relatively (warm California) undercurrent water which flows around the perimeter of the gyre. At lower (1.25-5 kHz) frequencies, the greater scattering strengths are found in the relatively cold water such is found in the upwelled subarctic water at the center of the gyre. This implies a significant change in the type of scatterers between these frequency domains. <i>Keywords:</i>					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL David G. Browning			22b. TELEPHONE (Include Area Code) (203)440-4173		22c. OFFICE SYMBOL 3331



**A RELATIONSHIP BETWEEN
OCEAN CIRCULATION AND VOLUME
REVERBERATION
IN THE
SUBARCTIC NORTHEAST PACIFIC OCEAN
(GULF OF ALASKA)**

**D.G. BROWNING
NEW LONDON LABORATORY
NAVAL UNDERWATER SYSTEMS CENTER
NEW LONDON, CT 06320**

**R.G. TURNER AND J.W. POWELL
DEFENCE RESEARCH ESTABLISHMENT PACIFIC
FMO, VICTORIA, B.C.
VOS IBO, CANADA**

L60112a


VIEWGRAPH 1

This paper presents an apparent relationship between ocean circulation and volume reverberation in the Subarctic Northeast Pacific Ocean.

-- Next viewgraph, please. --

Approved	
NTIS	✓
Library	
U.S. Navy	
Joint	
By	
Date	
Date	
A-1	





L60214a

VIEWGRAPH 2


In the North Pacific Ocean the principal circulation gyre extends from the equator to only 40° North latitude (corresponding to the coast of Northern California). This leaves a large Subarctic region to the North that has its own unique characteristics.

-- Next viewgraph, please. --



NORTH PACIFIC INTEGRATED SCATTERING STRENGTH LATITUDE DEPENDENCE (2.5-5.0 kHz BAND)

(J.A. SCRINGER AND R.G. TURNER, J ACOUST SOC AM 54(2), 483-493, (1973))



L60112d

VIEWGRAPH 3


Scrimger and Turner have previously shown a significant change in integrated scattering strength when transiting into the Subarctic region. This was a track from Vancouver to Hawaii and back into the Subarctic region.

-- Next viewgraph, please. --



NORTH PACIFIC COLUMN STRENGTH SPECTRA

(1.3) EFFECTIVE SWIM BLADDER RADIUS (cm)



(R.P. CHAPMAN et al, J ACOUSTIC SOC AM 56 (6), 1722 - 1734 (1974))

L60112c


VIEWGRAPH 4

Chapman, et al., as part of their epic circumnavigation of the western hemisphere, reported a change in the spectral content of column strength for two stations in this Subarctic region. Specifically, there appeared to be two independent mechanisms, one below 5 kilohertz and one (or more) above.

-- Next viewgraph, please. --



CIRCULATION SYSTEMS IN THE SUBARCTIC NORTH PACIFIC




L60214b


VIEWGRAPH 5

The circulation in the Subarctic Northeast Pacific is dominated by flow around the Alaskan gyre. Relatively warm water circulates around the perimeter while relatively cold water upwells in the center of the gyre.

-- Next viewgraph, please. --



September 1971 stations showing station numbers and positions. Open and closed circles indicate daytime and nighttime stations, respectively.*




April 1972 stations showing station numbers and positions. Open and closed circles indicate daytime and nighttime stations, respectively.*

L60214e

VIEWGRAPH 6

To quantify volume reverberation throughout the Northeast Subarctic Pacific (Gulf of Alaska), Turner conducted an extensive series of measurements for both summer and winter conditions. The standard measurement technique with explosive sources was used and data were reported for four octave frequency bands - 1.25 to 2.5 kHz, 2.5 to 5 kHz, 5 to 10 kHz, and 10 to 20 kHz. Night stations are shown by dark circles, day stations by light circles.

-- Next viewgraph, please. --



L80214g


VIEWGRAPH 7

Turner initially plotted the data as a function of latitude for all but one track. Typical results are shown here for the 10 to 20 kHz and 5 to 10 kHz bands. Rather than showing a simple trend the data have some unexpected changes.

-- Next viewgraph, please. --



CONTOURS OF THE AXIS DEPTH OF THE DEEP SOUND CHANNEL (SUMMER)



L60214c

S 81


VIEWGRAPH 8

Recently Powell, Chow, and Browning have shown that the depth of the deep sound channel axis in the Subarctic Northeast Pacific does not decrease with latitude but rather contours are concentric about the Alaskan gyre. The deepest axis depth is associated with the warm water around the perimeter, the shallowest with the cold water in the center of the gyre.

-- Next viewgraph, please. --



CONTOURS OF THE AXIS DEPTH OF THE DEEP SOUND CHANNEL (WINTER)



L60214d


VIEWGRAPH 9

This pattern intensifies under winter conditions with the axis reaching the surface in the center of the gyre.

-- Next viewgraph, please. --



COLUMN SCATTERING STRENGTH MEASUREMENTS AND CONTOURS FOR THE 10-20 kHz BAND



L60214h

VIEWGRAPH 10


It seemed logical to analyze Turner's data to see if a similar pattern developed. Here is the result for the highest frequency band (10 to 20 kilohertz), the solid contours are for summer, the dashed contours for winter. The highest column scattering strengths are associated with the warmer water around the perimeter of the gyre; here levels are generally lower under winter conditions.

The lowest levels are found in the colder water at the center of the gyre.

-- Next viewgraph, please. --



COLUMN SCATTERING STRENGTH MEASUREMENTS AND CONTOURS FOR THE 5-10 kHz BAND




VIEWGRAPH 11

A similar pattern is found for the next lower frequency band, 5 to 10 kilohertz. In all these figures, we are showing day stations only.

-- Next viewgraph, please. --




COLUMN SCATTERING STRENGTH MEASUREMENTS AND CONTOURS FOR THE 2.5-5 kHz BAND





COLUMN SCATTERING STRENGTH MEASUREMENTS AND CONTOURS FOR THE 1.25-2.5 kHz BAND



VIEWGRAPH 13

This pattern is confirmed by the lowest frequency band, 1.25 to 2.5 kilohertz. As Chapman's earlier work had indicated, we appear to have two distinct and independent frequency regimes with the demarcation being at 5 kilohertz.

-- Next viewgraph, please. --



CONCLUSIONS

SUBARCTIC NORTHEAST PACIFIC OCEAN (GULF OF ALASKA)

- **AT HIGHER FREQUENCIES (5-20 kHz)
GREATER VOLUME REVERBERATION FOUND
IN WARMER WATER AROUND NORTHERN
AND EASTERN PERIMETER OF ALASKAN GYRE**
- **AT LOWER FREQUENCIES (1.25-5 kHz)
GREATER VOLUME REVERBERATION FOUND
IN COLDER WATER UPWELLED IN THE
CENTER OF THE ALASKAN GYRE**
- **RESULTS SUGGEST THE POSSIBILITY OF TWO
DISTINCT SCATTERING REGIMES
AND MECHANISMS**

L60112b

VIEWGRAPH 14

We can summarize our results as follows:

- At higher frequencies (5 to 20 kilohertz) greater volume reverberation is found in warmer water around the perimeter of the Alaskan gyre.
- For lower frequencies (1.25 to 5 kilohertz) greater volume reverberation is found in relatively colder water.

These results suggest the possibility of two distinct scattering regimes and mechanisms.

